

Hospital Protocol RSD/CRPS Patient: Handle With Care!

Reflex Sympathetic Dystrophy(RSD) also known as Complex Regional Pain Syndrome (CRPS) is a chronic condition characterized by severe burning pain, pathological changes in bone and skin, excessive sweating, tissue swelling and extreme sensitivity to touch. People afflicted with RSD/CRPS are extraordinarily sensitive to certain stimuli, such as touch, movement, and injections.

Patient Tips

1. Bring a written copy of your medication regime. It is important to verify whether your medication regime can be taken care of solely through use of the hospital's pharmacy. Some medications may not be part of the hospital's formulary.
2. Avoid having ice applied to the RSD-affected limb
3. Instruct all hospital staff to always ask before touching you!
4. Request that the surgery protocols be adjusted so that the pre-op shave be done after anesthesia (catheter is in place for epidural or whatever). It feels like a lawnmower has been run over your legs when they shave without anesthesia.

Patient Room

1. Whenever possible patient should be in a quiet part of the hospital
2. In a semi-private room, patient should be in the second bed to avoid inadvertent bumping
3. A Zone-Air bed should be used (adjust mattress pressure to patient preference)
4. Heat and air conditioning should be well regulated
5. Standing orders should be issued for patients to have warm blankets
6. Foot Cradle to hold bed linens off body area (will diminish tactile stimuli)
7. Frequent linen changes may be necessary due to hyperhidrosis (increased sweating).
8. Allow family members to bring in clean sheet and pillowcases for patient. Hospital sheets are not soft and can irritate the skin.
9. Place a sign above bed designating affected limb

The Joint Commission on Accreditation of Healthcare Organizations has mandated the healthcare institutions that they accredit to assess and treat your pain. Pain is now to be assessed as the fifth vital sign. Patients have the right to demand pain control and to be pain free.

Procedures

1. Perform a phlebotomy on unaffected limb only
2. Use Pediatric needles (any trauma can cause the spread of RSD to a new site)
3. Warm Alcohol or Betadine wipes with warm running water on outside of package before opening package (these wipes can be very cold to the patient)
4. If PICA site is available, see if blood can be obtained from PIC instead of using vena puncture technique.

Blood Pressure and Pulse Rate

1. Use cuff on unaffected limb only
2. Use thigh cuff if both upper extremities are affected

Transport

1. Ask what kind of help the patient needs when transferring to a stretcher or wheelchair (simply touching arms or legs may cause hyperalgesia)
2. Use extreme care over bumps, such as elevator doorways

Feeding

1. Be careful not to touch patient extremities with bedside table

Patient Identification

1. Place red bracelet on unaffected limb
2. Place red dot sticker on patient chart



PO Box 502
Milford, CT 06460
Toll-free: 877-662-7737
info@rsds.org
www.rsds.org