

# Systematic mutation analysis of seven dystonia genes in complex regional pain syndrome with fixed dystonia

M. Florencia Gosso · Annetje M. de Rooij · Elisenda Alsina-Sanchis ·  
Jessica T. Kamphorst · Johan Marinus · Jacobus J. van Hilten ·  
Arn M. J. M. van den Maagdenberg

Received: 7 September 2009 / Revised: 24 November 2009 / Accepted: 9 December 2009  
© Springer-Verlag 2009

**Abstract** Complex regional pain syndrome type 1 (CRPS-1) is a chronic pain disorder that in some patients is associated with fixed dystonia. The pathogenesis of CRPS and its relation to dystonia remain poorly understood. Several genes (so-called DYT genes) identified in other causes of dystonia play a role in mechanisms that have been implicated in CRPS. Because different mutations in the same gene can result in diverse phenotypes, we sequenced all coding exons of the DYT1, DYT5a, DYT5b, DYT6, DYT11, DYT12, and DYT16 genes in 44 CRPS patients with fixed dystonia to investigate whether high-penetrant causal mutations play a role in CRPS. No such mutations were identified, indicating that these genes do not seem to play a major role in CRPS.

**Keywords** Complex regional pain syndrome · Fixed dystonia · DYT genes

## Introduction

Complex regional pain syndrome type 1 (CRPS-1) is a chronic pain condition that is commonly preceded by an injury to an arm or a leg, although spontaneous onset has been reported in 7% of the cases [13, 49]. Skin sensitivity, swelling, sweating and skin color and temperature changes are other features typical of the acute phase of the syndrome [34]. About 25% of CRPS patients also develop abnormal postures (fixed dystonia) of the affected extremity [22, 43]. Dystonia is a movement disorder in which twisting or repetitive movements or sustained postures are caused by involuntary, sustained muscle contractions [17].

The pathogenesis of CRPS and its relation to dystonia remain poorly understood. The identification of genes and signalling pathways that confer susceptibility to CRPS or dystonia in CRPS may therefore provide valuable insights on how host response mechanisms may turn aberrant in response to tissue injury. There is evidence to suggest that genetic factors may play a role in CRPS. The syndrome may cluster in families [1, 6, 11, 16, 19, 23, 49]. Although several genetic associations between CRPS and the HLA gene complex on the short arm of chromosome 6 (6p21.3) have been reported [12, 27, 33, 46, 48], to date no single causative gene has been identified. In contrast to CRPS, there have been many successes in identifying genes for primary dystonia (i.e. DYT1, DYT3), (for review see [36]) as well as genes involved in syndromes in which dystonia is part of a broader clinical spectrum (also referred to as dystonia plus syndromes) (i.e. DYT5a, DYT5b, DYT11, DYT12). Close to 20 chromosomal loci (so-called DYT loci) have been identified for primary dystonia and dystonia plus syndromes. For ten of them, the causative gene has been identified: DYT1 (*TorsinA* [39]), DYT3 (*TAF1* [26]),

---

M. F. Gosso and A. M. de Rooij contributed equally to this work.

**Electronic supplementary material** The online version of this article (doi:10.1007/s00415-009-5426-6) contains supplementary material, which is available to authorized users.

---

M. F. Gosso (✉) · E. Alsina-Sanchis · J. T. Kamphorst ·  
A. M. J. M. van den Maagdenberg  
Department of Human Genetics, Leiden University Medical  
Center, PO Box 9600, 2300 RC Leiden, The Netherlands  
e-mail: M.F.Gosso@lumc.nl

A. M. J. M. van den Maagdenberg  
e-mail: A.M.J.M.van\_den\_Maagdenberg@lumc.nl

A. M. de Rooij · J. Marinus · J. J. van Hilten ·  
A. M. J. M. van den Maagdenberg  
Department of Neurology, Leiden University Medical Center,  
Leiden, The Netherlands

DYT5a (*GCHI* [25]), DYT5b (*TH* [31]), DYT6 (*THAP1* [18]), DYT8 (*MRI* [30, 41]), DYT11 (*SGCE* [52]), DYT12 (*ATPIA3* [10]), DYT16 (*PRKRA* [7]), and DYT18 (*SLC2A1* [50]).

Dystonia in CRPS may spread to other extremities, leaving some patients severely disabled [47]. There are reasons to suspect a higher genetic load in CRPS patients with fixed dystonia. The age at onset of CRPS patients with dystonia is, on average, 11 years younger than that of CRPS patients without dystonia [47], which suggests a larger role for genetic factors in this more affected subgroup of patients. Notably, two studies that found a positive association with the HLA complex investigated CRPS patients with fixed dystonia [12, 46]. We hypothesized that dystonia genes may confer susceptibility to CRPS or dystonia in CRPS. To increase our chances of finding gene mutations, we sequenced all coding exons of those DYT genes with a known role in biological pathways that may potentially express features of CRPS.

## Patients and methods

### Patients

Forty-four patients who visited our clinic with an early onset of CRPS-1 (<40 years) and dystonia in at least one extremity were considered for inclusion in the study (Table 1). CRPS was diagnosed according to the criteria of the International Association of the Study of Pain (IASP) [34]. Fixed dystonia was diagnosed by a neurologist with expertise in movement disorders (JJvH). Patient clinical characteristics are given in Supplementary Table 1. The study was approved by the Medical Ethical Committee of the Leiden University Medical Center. All patients gave written informed consent before participation.

### DYT gene selection

To increase the chance of finding gene mutations, we investigated DYT genes with a role in biological pathways involved in disease mechanisms that are considered to play a role in CRPS, or disease mechanisms that potentially may express one or more features encountered in CRPS. Of course, additional scientific information on CRPS pathology and/or DYT gene function may become available in the future, in which case additional DYT genes should be selected for investigation.

Currently, multiple mechanisms, including increased oxidative stress [15, 45], aberrant inflammation [4, 24], apoptosis [37] and aberrant neuroplasticity [14, 32, 37], have been suggested as being involved in the pathogenesis of CRPS-I. Additionally, DYT genes that potentially may

**Table 1** Characteristics of patients with complex regional pain syndrome (CRPS) with fixed dystonia

Number of patients	44	
Percentage ( <i>N</i> ) of females	96%	(42)
Mean (SD) age at onset of CRPS (years)	26.5	(9.1)
Mean (SD) age at onset of dystonia (years)	27.0	(8.8)
Mean disease duration (SD) (years)	14.2	(7.1)
Family history of CRPS—percentage ( <i>N</i> )	27%	(12)
Preceding trauma—percentage ( <i>N</i> )		
Fracture	11%	(5)
Surgery	23%	(10)
Soft tissue	43%	(19)
Non-trauma	23%	(10)
Extremities affected by CRPS—percentage ( <i>N</i> )		
1	11%	(5)
2	32%	(14)
3	18%	(8)
4	39%	(17)
Extremities affected by dystonia—percentage ( <i>N</i> )		
1	23%	(10)
2	41%	(18)
3	14%	(6)
4	14%	(6)
First affected extremity—percentage ( <i>N</i> )		
Arm	52%	(23)
Leg	45%	(20)
Both	2%	(1)

Some patients had already received treatment before the start of the current study, therefore it was possible that at the time of evaluation the dystonia in these patient was no longer observed (*N* = 4)

*N* number, *SD* standard deviation

play a role in expressing phenotypic characteristics, such as pain and susceptibility to external triggers in initiating the phenotype, were included in this study. Hence, seven out of ten DYT genes (i.e., DYT1, DYT5a, DYT5b, DYT6, DYT11, DYT12, and DYT16) were selected for this study.

Our rationale for this selection was as follows:

- Mutations in *Torsin A* (DYT1) cause early-onset torsion, the most common and severe form of hereditary primary torsion dystonia [39]. Its gene product is implicated in the response of neurons to enhanced oxidative stress [28] and perhaps plasticity changes [20].
- *GCHI* (DYT5a), which codes for GTP cyclohydrolase 1, the rate-limiting enzyme in the biosynthesis of tetrahydrobiopterin (BH4) is a critical factor in neuropathic and inflammatory pain [44]. BH4 is also able to reduce the inflammatory response and oxidative stress [2]. Notably, a polymorphism in *GCHI* is associated with pain sensitivity [44].

- *TH* (DYT5b), which codes for tyrosine hydroxylase, the rate-limiting enzyme in the synthesis of the neurotransmitter dopamine [21], which is involved in modulation of pain perception. Not only do decreased levels of dopamine contribute to painful symptoms [51], but dopamine may also inhibit upregulation of cytokines and induce the production of anti-inflammatory mediators [3]. Finally, dopamine has a prolonged effect on neuroplasticity [29].
- Mutations in *THAP1* (DYT6), a member of a family of cellular factors responsible for regulation of endothelial cell proliferation [8], acting also as a proapoptotic factor [42], have been recently linked to primary forms of dystonia [18].
- *SGCE* (DYT11) gene encodes  $\epsilon$ -sarcoglycan, a protein in dopaminergic neurons in the substantia nigra and ventral tegmental area [9, 38] that seems to play a role in the synaptic function of the central nervous system [38].
- Mutations in *ATPIA3* (DYT12), which encodes the Na<sup>+</sup>/K<sup>+</sup>-ATPase  $\alpha$ 3 subunit can cause an increased cortical motor excitability and disruption of basal ganglia inhibitory control, leading to abrupt onset dystonia after physical or emotional trauma [5].
- *PRKRA* (DYT16), which codes for the protein kinase interferon-inducible double stranded RNA-dependent activator and plays an important role in response to extracellular stress and inflammatory cytokines interferon- $\gamma$  and TNF- $\alpha$  [7, 40].

We could not find a clear rationale for selecting the remaining three DYT genes (i.e., DYT3, DYT8, and DYT18).

#### Mutation analysis in selected DYT genes

For the genetic analysis, genomic DNA was isolated from peripheral blood cells according to a standard salting-out method [35]. All coding exons and directly adjacent intronic sequences of the *TorsinA*, *CGHI*, *TH*, *THAP1*, *SGCE*, *ATPIA3*, and *PRKRA* genes were analyzed for mutations using direct sequencing analysis. Exons were amplified by PCR using exon-specific primers sets (Supplementary Table 2). Reactions were performed in a 25- $\mu$ L reaction volume, containing 10 pmol of each primer, 1 $\times$  PCR buffer (3 mM Tris-HCl, 75 mM NH<sub>4</sub>SO<sub>4</sub>, 7.5 mM MgCl<sub>2</sub> with pH 8.5) (Invitrogen, Breda, The Netherlands), 3 mM dNTPs, 0.25 U AmpliTaq DNA polymerase (Applied Biosystems, Nieuwerkerk aan den IJssel, The Netherlands), and 50 ng of genomic DNA. PCR conditions were as follows: 3 min at 94°C, followed by 33 cycles of 30 s at 94°C, 30 s at 60°C, and 1 min at 72°C, and an additional extension step of 10 min at 72°C. Unincorporated dNTPs and primers were

removed by incubation at 37°C for 2 h with shrimp alkaline phosphatase (SAP) (USB Corporation, Cleveland, OH, USA) and exonuclease I (ExoI) (USB Corporation, Cleveland, OH, USA), followed by a deactivation step of 95°C for 15 min. For dideoxy sequencing, purified PCR product (15–25 ng) was used with 6 pmol forward or reverse primer in a final volume of 12  $\mu$ L. Sequencing reactions were run on an automated sequencer (ABI3730, Applied Biosystems, Nieuwerkerk aan den IJssel, The Netherlands). Sequence analysis was performed using ContigExpress software (a component of vector NTI Suite V9.0.0; Invitrogen, Leek, The Netherlands).

#### Results

Here we sequenced all coding exons and directly adjacent intronic sequences of the DYT1, DYT5a, DYT5b, DYT6, DYT11, DYT12, and DYT16 genes in 44 CRPS patients with fixed dystonia and an early-onset of disease (Table 1). The mean age at onset of CRPS in patients was 26.5 [standard deviation (SD) 9.1]. The mean age at onset of dystonia in patients was 27.0 (SD 8.8). In 23% ( $N = 10$ ) of patients, CRPS and dystonia developed without an initiating traumatic event. Some 89% ( $N = 39$ ) had two or more extremities with CRPS symptoms, whereas 77% ( $N = 34$ ) had two or more extremities with dystonia. In 27% ( $N = 12$ ) of the patients a positive family history of CRPS was present. No causal mutations were identified in any of the patients with CRPS and fixed dystonia.

#### Discussion

The pathogenesis of complex regional pain syndrome type 1 (CRPS-1) is poorly understood. Evidence for genetic factors, especially the involvement of the HLA complex on chromosome 6, in CRPS is increasing [12, 27, 33, 46, 48]. Despite these achievements, no single genetic factor has been identified in CRPS.

About 25% of CRPS patients develop movement disorders, of which dystonia is the most common one [22, 43, 49]. We hypothesized that those DYT genes that are responsible for monogenic forms of dystonias, could play a role conferring susceptibility to develop fixed dystonia in CRPS patients, and perhaps also CRPS itself.

Seven primary dystonia and dystonia plus syndromes genes (namely DYT1, DYT5a, DYT5b, DYT6, DYT11, DYT12, and DYT16) were selected because of their role in biological pathways (i.e., an abnormal response to oxidative stress, involvement in inflammation, apoptosis, and/or neuroplasticity) that also have been suggested to be involved in CRPS or in the expression of clinical

characteristics typical of the syndrome (e.g., pain, a triggering event). No disease-causing mutations were found in any of these genes tested in a fairly large group of CRPS patients with fixed dystonia and an early onset of the disease.

Although we did not find any evidence for the involvement of these DYT genes in CRPS with fixed dystonia, we cannot exclude the possibility that certain mutations were missed by our direct sequencing approach. Moreover, as we a priori cannot predict whether causal mutations in CRPS with fixed dystonia are loss- or gain-of-function mutations, the problem of missing mutations may particularly be true in the case in which causal mutations would be of the former category. Loss-of-function mutations that are located in the promoter region, or in other regulatory sequences affecting gene expression levels, would remain undetected with our mutation analysis approach. The same is true for causal deletions or insertions that interfere with any of the primer binding sites. Finally, the possibility that low-penetrant variants in DYT genes confer an increased risk for CRPS with fixed dystonia was not tested. In this respect, it is worth mentioning again that a polymorphism in *GCHI* has been associated with pain sensitivity [44]. Our current patient sample is too small to perform meaningful association studies aimed at identifying such low-penetrant gene variants.

A definite conclusion that can be drawn from the present study is that exonic mutations that cause dystonia in the DYT phenotypes belonging to the seven genes that were investigated are *not* present in CRPS with fixed dystonia. In that sense, we can conclude that the pathogenesis in the ‘DYT types’ of dystonia differs from that in CRPS or dystonia in CRPS. As some of the DYT genes have only recently been identified (i.e., *SCL2A1* in DYT18) and heavily studied, it still is not entirely clear whether these genes do not play a role in relevant pathways of CRPS with fixed dystonia. Therefore, as knowledge of CRPS and of the biology of DYT genes is increasing, other DYT genes may become a target for a study like the one performed here. There remains the possibility, of course, that none of the DYT genes play a role in CRPS with fixed dystonia.

**Acknowledgments** This study is part of Trauma RElated Neuronal Dysfunction (TREND), a Dutch Consortium that integrates research on epidemiology, assessment technology, pharmacotherapeutics, biomarkers and genetics on Complex Regional Pain Syndrome type 1. The consortium aims to develop concepts on disease mechanisms that occur in response to tissue injury, its assessment and treatment. TREND is supported by a government grant (BSIK03016).

**Conflict of interest statement** The authors declare that they have no conflict of interest.

## References

1. Albert J, Ott H (1983) Three brothers with algodystrophy of the hip. *Ann Rheum Dis* 42:421–424
2. Ali ZA, Bursill CA, Douglas G, McNeill E, Papaspyridonos M, Tatham AL, Bendall JK, Akhtar AM, Alp NJ, Greaves DR, Channon KM (2008) CCR2-mediated antiinflammatory effects of endothelial tetrahydrobiopterin inhibit vascular injury-induced accelerated atherosclerosis. *Circulation* 118:S71–S77
3. Beck G, Brinkkoetter P, Hanusch C, Schulte J, van Ackern K, van der Woude FJ, Yard BA (2004) Clinical review: immunomodulatory effects of dopamine in general inflammation. *Crit Care* 8:485–491
4. Birklein F, Schmelz M (2008) Neuropeptides, neurogenic inflammation and complex regional pain syndrome (CRPS). *Neurosci Lett* 437:199–202
5. Blanco-Arias P, Einholm AP, Mamsa H, Concheiro C, Gutierrez-de-Teran H, Romero J, Toustrup-Jensen MS, Carracedo A, Jen JC, Vilsen B, Sobrido MJ (2009) A C-terminal mutation of ATP1A3 underscores the crucial role of sodium affinity in the pathophysiology of rapid-onset dystonia-parkinsonism. *Hum Mol Genet* 18:2370–2377
6. Brusca IC, Beltran Audera CH, Jimenez ZF (2004) Polytopic and recurrent reflex sympathetic dystrophy in lower limbs in two siblings. *An Med Int* 21:183–184
7. Camargos S, Scholz S, Simon-Sanchez J, Paisan-Ruiz C, Lewis P, Hernandez D, Ding J, Gibbs JR, Cookson MR, Bras J, Guerreiro R, Oliveira CR, Lees A, Hardy J, Cardoso F, Singleton AB (2008) DYT16, a novel young-onset dystonia-parkinsonism disorder: identification of a segregating mutation in the stress-response protein PRKRA. *Lancet Neurol* 7:207–215
8. Cayrol C, Lacroix C, Mathe C, Ecochard V, Ceribelli M, Loreau E, Lazar V, Dessen P, Mantovani R, Aguilar L, Girard JP (2007) The THAP-zinc finger protein THAP1 regulates endothelial cell proliferation through modulation of pRB/E2F cell-cycle target genes. *Blood* 109:584–594
9. Chan P, Gonzalez-Maeso J, Ruf F, Bishop DF, Hof PR, Sealton SC (2005) Epsilon-sarcoglycan immunoreactivity and mRNA expression in mouse brain. *J Comp Neurol* 482:50–73
10. de Carvalho Aguiar P, Sweadner KJ, Penniston JT, Zarella J, Liu L, Caton M, Linazasoro G, Borg M, Tijssen MA, Bressman SB, Dobyns WB, Brashear A, Ozelius LJ (2004) Mutations in the Na<sup>+</sup>/K<sup>+</sup>-ATPase alpha3 gene ATP1A3 are associated with rapid-onset dystonia parkinsonism. *Neuron* 43:169–175
11. de Rooij AM, de Mos M, Sturkenboom MC, Marinus J, van den Maagdenberg AM, van Hilten JJ (2009a) Familial occurrence of complex regional pain syndrome. *Eur J Pain* 13:171–177
12. de Rooij AM, Florencia Gosso M, Haasnoot GW, Marinus J, Verduijn W, Claas FH, van den Maagdenberg AM, van Hilten JJ (2009b) HLA-B62 and HLA-DQ8 are associated with complex regional pain syndrome with fixed dystonia. *Pain* 145:82–85
13. de Rooij AM, Perez RS, Huygen FJ, Eijs FV, Kleef MV, Bauer MC, van Hilten JJ, Marinus J (2009c) Spontaneous onset of complex regional pain syndrome. *Eur J Pain* PMID 19793666
14. Drummond PD, Finch PM (2006) Sensory changes in the forehead of patients with complex regional pain syndrome. *Pain* 123:83–89
15. Eisenberg E, Shtahl S, Geller R, Reznick AZ, Sharf O, Ravbinovich M, Erenreich A, Nagler RM (2008) Serum and salivary oxidative analysis in complex regional pain syndrome. *Pain* 138:226–232
16. Erdmann MW, Wynn-Jones CH (1992) ‘Familial’ reflex sympathetic dystrophy syndrome and amputation. *Injury* 23:136–138
17. Fahn S (1988) Concept and classification of dystonia. *Adv Neurol* 50:1–8

18. Fuchs T, Gavarini S, Saunders-Pullman R, Raymond D, Ehrlich ME, Bressman SB, Ozelius LJ (2009) Mutations in the THAP1 gene are responsible for DYT6 primary torsion dystonia. *Nat Genet* 41:286–288
19. Galer BS, Henderson J, Perander J, Jensen MP (2000) Course of symptoms and quality of life measurement in complex regional pain syndrome: a pilot survey. *J Pain Symptom Manage* 20:286–292
20. Granata A, Schiavo G, Warner TT (2009) TorsinA and dystonia: from nuclear envelope to synapse. *J Neurochem* 109:1596–1609
21. Grima B, Lamouroux A, Boni C, Julien JF, Javoy-Agid F, Mallet J (1987) A single human gene encoding multiple tyrosine hydroxylases with different predicted functional characteristics. *Nature* 326:707–711
22. Harden RN, Bruhl S, Galer BS, Saltz S, Bertram M, Backonja M, Gayles R, Rudin N, Bhugra MK, Stanton-Hicks M (1999) Complex regional pain syndrome: are the IASP diagnostic criteria valid and sufficiently comprehensive? *Pain* 83:211–219
23. Huhne K, Leis S, Schmelz M, Rautenstrauss B, Birklein F (2004) A polymorphic locus in the intron 16 of the human angiotensin-converting enzyme (ACE) gene is not correlated with complex regional pain syndrome I (CRPS I). *Eur J Pain* 8:221–225
24. Huygen FJ, De Bruijn AG, De Bruin MT, Groeneweg JG, Klein J, Zijlstra FJ (2002) Evidence for local inflammation in complex regional pain syndrome type I. *Mediators Inflamm* 11:47–51
25. Ichinose H, Ohye T, Takahashi E, Seki N, Hori T, Segawa M, Nomura Y, Endo K, Tanaka H, Tsuji S et al (1994) Hereditary progressive dystonia with marked diurnal fluctuation caused by mutations in the GTP cyclohydrolase I gene. *Nat Genet* 8:236–242
26. Kaji R, Goto S, Tamiya G, Ando S, Makino S, Lee LV (2005) Molecular dissection and anatomical basis of dystonia: X-linked recessive dystonia-parkinsonism (DYT3). *J Med Invest* 52(suppl):280–283
27. Kemler MA, van de Vusse AC, van den Berg-Loonen EM, Barendse GA, van Kleef M, Weber WE (1999) HLA-DQ1 associated with reflex sympathetic dystrophy. *Neurology* 53:1350–1351
28. Kuner R, Teismann P, Trutzel A, Naim J, Richter A, Schmidt N, Bach A, Ferger B, Schneider A (2004) TorsinA, the gene linked to early-onset dystonia, is upregulated by the dopaminergic toxin MPTP in mice. *Neurosci Lett* 355:126–130
29. Kuo MF, Paulus W, Nitsche MA (2008) Boosting focally-induced brain plasticity by dopamine. *Cereb Cortex* 18:648–651
30. Lee HY, Xu Y, Huang Y, Ahn AH, Auburger GW, Pandolfo M, Kwiecinski H, Grimes DA, Lang AE, Nielsen JE, Averyanov Y, Servidei S, Friedman A, Van Bogaert P, Abramowicz MJ, Bruno MK, Sorensen BF, Tang L, Fu YH, Ptacek LJ (2004) The gene for paroxysmal non-kinesigenic dyskinesia encodes an enzyme in a stress response pathway. *Hum Mol Genet* 13:3161–3170
31. Ludecke B, Dworniczak B, Bartholome K (1995) A point mutation in the tyrosine hydroxylase gene associated with Segawa's syndrome. *Hum Genet* 95:123–125
32. Maihofner C, Handwerker HO, Neundorfer B, Birklein F (2003) Patterns of cortical reorganization in complex regional pain syndrome. *Neurology* 61:1707–1715
33. Mailis A, Wade J (1994) Profile of Caucasian women with possible genetic predisposition to reflex sympathetic dystrophy: a pilot study. *Clin J Pain* 10:210–217
34. Merskey H, Bogduk N (1994) Complex Regional Pain Syndrome, Type I. In: Merskey H, Bogduk N (eds) *Classification of chronic pain: descriptions of chronic pain syndromes and definitions of pain terms*. IASP Press, Seattle, pp 41–42
35. Miller SA, Dykes DD, Polesky HF (1988) A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 16:1215
36. Muller U (2009) The monogenic primary dystonias. *Brain* 132:2005–2025
37. Navarro X, Vivo M, Valero-Cabre A (2007) Neural plasticity after peripheral nerve injury and regeneration. *Prog Neurobiol* 82:163–201
38. Nishiyama A, Endo T, Takeda S, Imamura M (2004) Identification and characterization of epsilon-sarcoglycans in the central nervous system. *Brain Res Mol Brain Res* 125:1–12
39. Ozelius LJ, Hewett JW, Page CE, Bressman SB, Kramer PL, Shalish K, de Leon D, Brin MF, Raymond D, Corey DP, Fahn S, Risch NJ, Buckler AJ, Gusella JF, Breakefield XO (1997) The early-onset torsion dystonia gene (DYT1) encodes an ATP-binding protein. *Nat Genet* 17:40–48
40. Patel CV, Handy I, Goldsmith T, Patel RC (2000) PACT, a stress-modulated cellular activator of interferon-induced double-stranded RNA-activated protein kinase, PKR. *J Biol Chem* 275:37993–37998
41. Rainier S, Thomas D, Tokarz D, Ming L, Bui M, Plein E, Zhao X, Lemons R, Albin R, Delaney C, Alvarado D, Fink JK (2004) Myofibrillogenesis regulator 1 gene mutations cause paroxysmal dystonic choreoathetosis. *Arch Neurol* 61:1025–1029
42. Roussigne M, Cayrol C, Clouaire T, Amalric F, Girard JP (2003) THAP1 is a nuclear proapoptotic factor that links prostatic apoptosis-response-4 (Par-4) to PML nuclear bodies. *Oncogene* 22:2432–2442
43. Schwartzman RJ, Kerrigan J (1990) The movement disorder of reflex sympathetic dystrophy. *Neurology* 40:57–61
44. Tegeer I, Costigan M, Griffin RS, Abele A, Belfer I, Schmidt H, Ehnert C, Nejm J, Marian C, Scholz J, Wu T, Allchome A, Diatchenko L, Binshtok AM, Goldman D, Adolph J, Sama S, Atlas SJ, Carlezon WA, Parsegian A, Lotsch J, Fillingim RB, Maixner W, Geisslinger G, Max MB, Woolf CJ (2006) GTP cyclohydrolase and tetrahydrobiopterin regulate pain sensitivity and persistence. *Nat Med* 12:1269–1277
45. van der Laan L, ter Laak HJ, Gabreels-Festen A, Gabreels F, Goris RJ (1998) Complex regional pain syndrome type I (RSD): pathology of skeletal muscle and peripheral nerve. *Neurology* 51:20–25
46. van Hilten JJ, van de Beek WJ, Roep BO (2000) Multifocal or generalized tonic dystonia of complex regional pain syndrome: a distinct clinical entity associated with HLA-DR13. *Ann Neurol* 48:113–116
47. van Rijn MA, Marinus J, Putter H, van Hilten JJ (2007) Onset and progression of dystonia in complex regional pain syndrome. *Pain* 130:287–293
48. Vaneker M, van der Laan L, Allebes WA, Gorski RJ (2002) Genetic factors associated with complex regional pain syndrome 1: HLA DRB and TNF alpha promoter gene polymorphism. *Disabil Med* 2:69–74
49. Veldman PH, Reynen HM, Arntz IE, Goris RJ (1993) Signs and symptoms of reflex sympathetic dystrophy: prospective study of 829 patients. *Lancet* 342:1012–1016
50. Weber YG, Storch A, Wuttke TV, Brockmann K, Kempfle J, Maljevic S, Margari L, Kamm C, Schneider SA, Huber SM, Pekrun A, Roebing R, Seeböhm G, Koka S, Lang C, Kraft E, Blazevic D, Salvo-Vargas A, Fauler M, Mottaghy FM, Munchau A, Edwards MJ, Presicci A, Margari F, Gasser T, Lang F, Bhatia KP, Lehmann-Horn F, Lerche H (2008) GLUT1 mutations are a cause of paroxysmal exertion-induced dyskinesias and induce hemolytic anemia by a cation leak. *J Clin Invest* 118:2157–2168
51. Wood PB (2008) Role of central dopamine in pain and analgesia. *Expert Rev Neurother* 8:781–797
52. Zimprich A, Grabowski M, Asmus F, Naumann M, Berg D, Bertram M, Scheidtman K, Kern P, Winkelmann J, Müller-Myhök B, Riedel L, Bauer M, Müller T, Castro M, Meitinger T, Strom TM, Gasser T (2001) Mutations in the gene encoding epsilon-sarcoglycan cause myoclonus-dystonia syndrome. *Nat Genet* 29:66–69