

REVIEW

Reflex sympathetic dystrophy—a complex regional pain syndrome

L. TURNER-STOKES*

Herbert Dunhill Chair in Rehabilitation, King's College London and Director, Regional Rehabilitation Unit, Northwick Park Hospital

Accepted for publication: January 2002

Abstract

Reflex sympathetic dystrophy (RSD) is a complex and poorly-understood condition characterized by: (a) pain and altered sensation; (b) motor disturbance and soft tissue change; (c) vasomotor and autonomic changes; and (d) psychosocial disturbance. Neurological symptoms typically do not conform to any particular pattern of nerve damage. Many different names have been ascribed to this condition and most recently the term 'complex regional pain syndrome' has been coined to emphasize the complex interaction of somatic, psychological and behavioural factors. Diagnostic criteria have been proposed by the International Association for the Study of Pain, but are still subject to debate. This review article describes the clinical features which may present as part of the condition, and the patho-physiology and pre-disposing factors so far identified. The evidence for effectiveness of different interventions is presented and a treatment approach outlined for inter-disciplinary management. While RSD is traditionally associated with pain in the extremities, the possibility is raised that the same process may underlie chronic pain syndromes affecting more central structures, such as testicular or pelvic pain.

Introduction

Reflex sympathetic dystrophy (RSD) is a complex and often poorly-understood condition characterized by:

- (a) pain and altered sensation;
- (b) motor disturbance and soft tissue change;
- (b) vasomotor and autonomic changes; and
- (b) psychosocial disturbance.

RSD may affect either upper or lower limb, may be localized, or may involve the whole extremity. Rarely

it affects more than one limb.^{1,2} Neurological symptoms typically do not conform to any particular pattern of nerve damage.

This review article represents an assimilation of the literature to describe the condition, its diagnosis and an approach to rehabilitation.

Terminology

A number of different names have been given to this condition including:

- Reflex sympathetic dystrophy (RSD);
- Algodystrophy;
- Sudek's atrophy;
- Shoulder-hand syndrome (SHS);
- Sympathetically-maintained pain (SMP);
- Complex regional pain syndrome (CRPS); and
- Causalgia.

The multiple terms are often confusing and make literature searching difficult. Some are applied to specific subgroups (e.g. Sudek's atrophy which describes the condition of RSD in the hand following Colles fracture or causalgia which derives from the Greek '*causos*' or 'heat' and describes the burning pain sometimes associated with neuropathic damage). Others (e.g. Complex regional pain syndrome) attempt to describe the general condition.

Until recently 'Reflex sympathetic dystrophy' was the most popular term. The reflex nature is unclear, however and the role of sympathetic system not well understood, so the term 'Complex regional pain syndrome' has been coined to emphasize the complex interaction of somatic, psychological and behavioural factors and the regional (i.e. non-localized) distribution of symptoms.³ The International Association for the Study of Pain (IASP) has recently attempted to develop criteria to improve the

* Address for correspondence: Regional Rehabilitation Unit, Northwick Park Hospital, Watford Road, Harrow, Middlesex, HA1 3UJ, UK; e-mail: lynne.turner-stokes@dial.pipex.com

diagnosis of CRPS. However, these have not been fully validated and need further work.

At the current time, 'Reflex sympathetic dystrophy' is the term more widely used in the literature and recognized by clinicians other than pain specialists. It is not entirely clear that all which is described in that huge literature on RSD would conform to the criteria for CRPS. This review leans heavily on the RSD literature, so that term will be used for the current purpose.

Clinical features

The main clinical features of RSD are listed in table 1.³⁻⁵ The pattern of presentation may alter somewhat with time and the clinical course is commonly divided into three stages:^{6, 7}

- (1) acute or hyperaemic;
- (2) ischaemic; and
- (3) atrophic.

In the early stages, vasomotor changes may be very prominent with acute pain, swelling, colour change and a 'hot' bone scan. In the later stages, atrophic changes predominate with pallor, contracture, and osteopenia, which may result in part from learned disuse.³

Pathophysiology

The often-striking response of RSD to sympathetic blockade together with signs of autonomic dysfunction supports the notion that it represents an exaggerated sympathetic response to post-traumatic inflammatory responses, rather than the consequence of sympathetic damage.⁸ The concept of sympathetically-maintained pain is supported by human and animal experiments, which demonstrate that catecholamines can activate primary afferent nociceptors, together with the fact that sympathetic blockade is sometimes effective in relieving symptoms. However, the complex interaction of psychological factors and failure to demonstrate increased levels of catecholamines and related factors peripherally⁹⁻¹¹ suggests that the processing problem is central rather than peripheral.³

What triggers RSD?

In 95% of cases there is some history of trauma or surgery.³ However, there is little relationship between the severity of the condition and of the original insult. Typically it follows minor injury or surgery.¹²

Table 1 Clinical features of reflex sympathetic dystrophy

Pain and altered sensation

- (1) Severe and very unpleasant pain
 - often graphically described as 'knife-life' or 'burning'
- (2) Altered sensation with
 - numbness
 - paraesthesia (pins and needles)
 - hyperalgesia (increased sensitivity to painful stimuli)
 - allodynia (normal stimuli perceived as pain)

Motor disturbance

- (1) Spasms—which may take the form of
 - dystonic posturing⁴
 - myoclonic type jerks at night—otherwise known as 'restless legs'⁵
- (2) Restriction of joint movement
 - usually limitation of flexion
 - joint tenderness/effusion
 - soft tissue contractures—develop within a relatively short space of time

Vasomotor/autonomic changes

- Variable skin colour changes from pallor to redness or purple, blotchy discoloration (this may have little relation to the ambient temperature, peripheral pulses are usually maintained)
- Temperature change—may be either warmer or colder than the unaffected limb
- Swelling and oedema—in the affected extremity
- Sweating change—either decrease or increase
- Trophic changes (shiny skin with hair loss)

Psychosocial

A complex interaction of psychological and behavioural factors, which may include

- enhanced disability behaviour—reflecting a greater level of dependence than would be explainable by the physical condition alone
- depression
- anger, distress, hostility—not infrequently there may be litigation underway
- physical and vocational inactivity—either preceding or since the onset of symptoms

RSD in the upper limb is reported following carpal tunnel release (whether performed endoscopically^{13, 14} or operatively¹⁵), Dupuytren's repair^{16, 17} and other tendon release procedures.¹⁸

RSD in the lower limb is a recognized complication of knee surgery—including tibial osteotomy^{19, 20} and arthroplasty²¹ where it is associated with a poor prognosis. It is also reported following crush injury to the foot,²² ankle arthrodesis,²³ amputation²⁴ and hip arthroplasty.²⁵

It is well-described following minor interventions such as arthroscopy, in which there is only minimal trauma.^{26, 27} One multi-centre study quotes a frequency of 2.3% following knee arthroscopy.²⁸ It is also reported following ankle arthroscopy^{29, 30} albeit less frequently.

In other reports, RSD has been associated with stroke,^{31, 32} mastectomy,³³ pregnancy³⁴ and osteogenesis imperfecta.³⁵

Predisposing factors

Certain psychological features are reported in association with RSD although it is sometimes difficult to separate cause and effect.

- Patients with industrial injuries have a higher rate of RSD for procedures such as arthroscopy,³⁶ and a number of studies report high frequencies where 'worker's compensation' insurance is involved.¹²
- There is a strong association with inactivity (including jobless persons and bankrupt self-employed people).³⁷
- Patients with RSD report more significant life events in the year preceding onset.^{38–40}
- Positive associations are also reported with a psychological history:⁴¹
 - (a) depression;^{37, 42–44}
 - (c) personality disorders;^{38, 45}
 - (c) psycho-neuroticism;³⁸
 - (c) hypochondriasis and hysteria;⁴⁶
 - (c) expressed helplessness;⁴⁷ and
 - (c) 'pain prone-ness'⁴⁸ or chronic pain.⁴⁹

RSD is frequently reported in association with an exaggeration of disability or signs, which may vary from simple functional overlay to actual malingering.^{50, 51} Results of treatment in compensation cases with pain dysfunction may be poor.⁵²

Diagnosis

The mainstay of diagnosis relies on the presence of clinical features appropriate to the stage of the condition. There have been no clearly defined criteria for the diagnosis of RSD. However, the IASP has recently developed a new set of diagnostic criteria for CRPS types I (without nerve damage) and II (with nerve damage). The criteria specifically avoid mention of the sympathetic nervous system or responsiveness to blockade. They include:

- Presence of an initiating noxious event or cause of immobilization.
- Continuing pain, allodynia or hyperalgesia disproportionate to the precipitating event.
- Evidence at some time of: oedema; changes in skin blood flow; sweating.
- No other condition which can account for the pain or dysfunction.

The criteria were developed by consensus and have not yet been fully validated. A multicentre study⁵³ to investigate validity, criticized the criteria which failed to include the motor and atrophic changes. The authors have proposed an experimental revision providing more stringent criteria for research purposes. These are:

- Continuing pain, allodynia or hyperalgesia disproportionate to the precipitating event.
- Symptoms in all four of the following areas:
 - (a) pain/sensitivity (hyperalgesia or allodynia);
 - (d) vasomotor—assymetry of skin temperature or colour;
 - (d) sudomotor/fluid—oedema or sweating change; and
 - (d) motor/dystrophic—reduced range of movement, motor patterns or atrophic change; and
- Signs in at least 2 of the four areas.

Both sets of criteria fail to describe psycho-social issues adequately and more work is clearly needed before either is fully accepted.³

INVESTIGATIONS

Standard X-rays typically demonstrate patchy demineralization in the affected part, which may be the result of disuse. Three-phase bone scintigraphy shows increased activity in both the blood pool and static phases.⁵⁴

A comparison of X-ray and bone scintigraphy⁵⁵ revealed that bone scans are more sensitive than X-rays, especially in the early phase of the condition. Three-phase scans are not necessary, since changes are equally well seen in the static phase. Bone scans appear to be sensitive to treatment and may therefore be used both for initial diagnosis and for monitoring response to treatment.

MRI has more recently gained recognition in diagnosis,^{56, 57} especially where radiology is contra-indicated, e.g. in pregnancy.³⁴ The following features are described:

- periarticular marrow oedema—often involving more than one bone;
- soft tissue swelling;
- joint effusions;⁵⁸ and
- a subchondral band of low T1-weighted intensity is a rare finding of questionable significance.

Infra-red telethermography is reported to be sensitive and specific⁵⁹ but is rarely used in clinical practice. Dexa scans are not useful for diagnosis, but may have a role in monitoring progress.⁶⁰

Management

If prevention is the best form of treatment, attention to analgesia and early mobilization post-operative may help to avoid the problem in the first place.^{61, 62} Peri-operative guanethidine blockade is not shown to have any protective effect, however.⁶³

Once developed, early recognition and proactive treatment are generally acknowledged to be associated with the best outcomes.⁶⁴ In the acute phase, sympathetic blockade can be strikingly effective, if short-lived, but may provide a useful window of pain-relief in which to encourage exercise and physiotherapy. Active physiotherapy is shown not only to be effective, but cost-effective.^{65–67} Treatment in the second or third stages of the disease is associated with a poorer outcome.⁶⁸

MEDICAL/SURGICAL INTERVENTIONS

A number of medical interventions have been tried with varying success (see table 2). The evidence for effectiveness has been reviewed by Kingery.⁶⁹

It is recognized, however, that medical management alone is insufficient to sustain benefit.⁸³ Sympathetic blockade can provide effective relief of symptoms in the short-term, but the effects are not long-lasting and symptoms inevitably recur. Permanent destruction of sympathetic nerves is currently considered inadvisable.⁸⁴ Here, too, symptoms recur, but now cannot even be blocked.³

There are relatively few properly controlled trials of these techniques, and this is a condition where the placebo effect may be expected to be strong. Low-tech treat-

Table 2 Medical/surgical interventions reported to be of benefit in RSD

Sympathetic blockade

- by intravenous injection with various agents⁷⁰
- by infiltration of the sympathetic plexus with local anaesthetic⁷¹

Destruction of the local sympathetic supply

- by dorsal or lumbar sympathectomy^{72–74}
- by radiofrequency lumbar sympathectomy⁷⁵

*Bisphosphonates*⁷⁶

*Calcitonin*⁷⁷

*Intra-thecal morphine*⁷⁸

Corticosteroids, given

- intravenously⁷⁹
- intra-muscularly⁸⁰
- orally³²

Spinal cord stimulation^{81,82}

ments such as sympathetic block or analgesia may provide a useful window of pain relief during which to win the patients' confidence and engage them in a rehabilitation programme.⁸⁵ Hi-tech treatments such as spinal cord stimulation should not be applied, other than as part of research trials, until their benefit and cost-effectiveness are proven.³

Inter-disciplinary approach to rehabilitation

In this complex condition, the longer symptoms and behaviours persist unchecked, the greater the interaction of psychosocial factors, and the worse the prognosis. A co-ordinated inter-disciplinary approach is needed.⁸⁶ Psychological support is reported to have a strongly positive effect on treatment outcomes and a cognitive behavioural approach with clearly functional goals and self-management techniques are recommended to maintain long-term independence.⁸⁷

TREATMENT GOALS INCLUDE:³

- Reduction of pain—if possible;
- reduction of pain behaviour;
- reduction of inappropriate use of health services—many patients go from doctor to doctor in search of a cure;
- increased function of the affected limb;
- increased quality of life and self-esteem; and
- improved psycho-social functioning and relationships with family/friends.

PROGRAMME STAGES³

To understand rehabilitation of the patient with RSD it is necessary to understand the sequence of events which occurs during its development (see figure 1). The critical stages of the programme are:^{87, 88}

Engage the patient

First of all it is necessary to engage the patient—they need to understand that the key to getting better is to use the limb, despite their experience that to do so hurts. Medical interventions can provide a window of pain relief during which the exercise programme gets underway.

Desensitization

Subject the patient to regular sensory stimulation, gradually building up both duration and intensity of stimulation. The rate of increase should be under the




Figure 1 Complex psychosocial/behavioural interaction deterioration in RSD

patient's control, according to clear goals which they are encouraged to set for themselves.

Flexibility

Physical techniques are applied to control oedema, increased range of movement and build up muscle strength with isometric exercises. Initially the programme of stretches and exercises may be passive

but, as soon as possible, responsibility is passed over to the patient to do their own programme of stretches and exercises, again with a clear set of goals.

Postural balance

Encourage the patient onto their feet for standing and balance control exercises (in lower limb RSD), aerobic conditioning and isotonic exercises.

Movement

Goals are now extended towards regaining normal movement patterns of dynamic function such as walking or dextrous movements. Mirrors and biofeedback techniques may be useful to increase awareness of undesired movement patterns and encourage the patients to self-monitor and correct them.

Vocational rehabilitation

Finally, once the best possible function is achieved, patients are followed through to apply that function in the workplace, if necessary modifying the job or work environment to enable productive activity. This is particularly important, since inactivity is seen to be a major contributor to the occurrence of RSD. If this stage is ignored, relapse is more likely. If productive work is impossible, efforts should focus on leisure and social activities instead.

Psycho-social/behavioural rehabilitation

Also crucial to the success of the rehabilitation programme is education of the patient and their families about the nature of the condition and how their behaviours can either perpetuate or improve it. The psycho-social issue must not be swept under the carpet. The patient needs to understand and accept the part played by these factors, how they affect their symptoms and what they can do about it.

They need to understand that theirs is a normal reaction to abnormal circumstances. They and their families need to be taught techniques for stress management, coping strategies, pacing and variation of activities, and reduced pain behaviour. Their families often need permission to withdraw and let the patient do tasks for themselves.

Other specific interventions

- Depression: if the patient is considered to be depressed, this should be treated with SSRIs or tricyclic anti-depressants. Many recommend the use of amitriptyline because of its supposed specific action on pain, but there is no hard research evidence to support this view.³
- Law-suits: if the patient is engaged in a law-suit, this should be facilitated to its conclusion as soon as possible. Litigation fuels anger, bitterness, stress and increases sympathetic arousal—rehabilitation is unlikely to be successful until litigation is completed one way or the other.³

RSD affecting other parts of the body

RSD has traditionally been thought of as affecting the limbs only. However, several authors have questioned whether there may be proximal manifestations of the condition.

Muller and colleagues⁸⁹ have recently suggested that frozen shoulder may be a manifestation of RSD and Julian and colleagues⁹⁰ have asked the same question with regard to vulvar vestibulitis.

Pelvic pain has frequently presented the medical profession with a diagnostic and management conundrum. In some cases there is a clearly identifiable cause such as endometriosis, or chronic pelvic inflammatory disease, but in some there is no clearly identifiable cause despite intensive investigation. Such patients quite frequently demonstrate the following features in common with RSD:

- a disproportionate level of distress and psychosocial dysfunction;⁹¹
- an initiating surgical procedure—often minor,⁹² and around which there may be litigation; and
- a previous psychological background.⁹³

It now seems likely that chronic regional pain syndromes may account for at least some cases of pelvic pain, and possibly also some cases of testicular or chronic abdominal pain. In some cases there may also be a sympathetically-maintained element to the pain and this may account for the reported success of testicular denervation (which includes the sympathetic supply) in some cases of intractable testicular pain.^{94,95} Either way, since effective treatment options are lacking, and since the condition may escalate into psycho-social breakdown and catastrophe if not handled effectively, a similar process interdisciplinary rehabilitation with strong cognitive/behavioural elements would appear to be a logical and sensible approach.

Acknowledgements

This paper represents an assimilation of the current literature and draws on the proceedings of a session entitled 'Complex regional pain syndrome: a fresh look at a difficult problem' at the 62nd Annual Assembly of American Academy of Physical Medicine and Rehabilitation, San Francisco, November 2000³, with thanks to the presenters of the session. Financial support for preparation of the article was kindly provided by the Luff Foundation.

References

- 1 Schiffenbauer J, Fagien M. Reflex sympathetic dystrophy involving multiple extremities. *Journal of Rheumatology* 1993; **20**(1): 165–169.
- 2 Teasell RW, Potter P, Moulin D. Reflex sympathetic dystrophy involving three limbs: a case study. *Archives of Physical Medicine & Rehabilitation* 1994; **75**(9): 1008–1010.
- 3 Bruehl S, Harden RN, Sorrell P. Complex regional pain syndromes, a fresh look at a difficult problem. 62nd Annual Assembly of the American Academy of Physical Medicine and Rehabilitation. San Francisco, 2000 (Conference proceedings).
- 4 Robberecht W, Van Hees J, Adriaensen H, Carton H. Painful muscle spasms complicating algodystrophy: central or peripheral disease? *Journal of Neurology, Neurosurgery and Psychiatry* 1988; **51**(4): 563–567.
- 5 DeWolfe VG. Iatrogenic and functional leg pain. *Geriatrics* 1973; **28**(9): 60–62.
- 6 Driessen JJ, van der Werken C, Nicolai JP, Crul JF. Clinical effects of regional intravenous guanethidine (Ismelin) in reflex sympathetic dystrophy. *Acta Anaesthesiologica Scandinavica* 1983; **27**(6): 505–509.
- 7 Kurvers HA, Jacobs MJ, Beuk RJ, van den Wildenburg FA, Kitslaar PJ, Slaaf DW *et al.* The influence of local skin heating and reactive hyperaemia on skin blood flow abnormalities in patients with reflex sympathetic dystrophy (RSD). *European Journal of Clinical Investigation* 1995; **25**(5): 346–352.
- 8 Baron R, Levine JD, Fields HL. Causalgia and reflex sympathetic dystrophy: does the sympathetic nervous system contribute to the generation of pain? *Muscle & Nerve* 1999; **22**(6): 678–695.
- 9 Drummond PD, Finch PM, Edvinsson L, Goadsby PJ. Plasma neuropeptide Y in the symptomatic limb of patients with causalgic pain. *Clinical Autonomic Research* 1994; **4**(3): 113–116.
- 10 Drummond PD, Finch PM, Smythe GA. Reflex sympathetic dystrophy: the significance of differing plasma catecholamine concentrations in affected and unaffected limbs. *Brain* 1991; **114**(pt 5): 2025–2036.
- 11 Goldstein DS, Tack C, Li ST. Sympathetic innervation and function in reflex sympathetic dystrophy. *Annals of Neurology* 2000; **48**(1): 49–59.
- 12 Finsterbush A, Frankl U, Mann G, Lowe J. Reflex sympathetic dystrophy of the patellofemoral joint. *Orthopaedic Review* 1991; **20**(10): 877–885.
- 13 Slattery PG. Endoscopic carpal tunnel release. Use of the modified Chow technique in 215 cases (see comments). *Medical Journal of Australia* 1994; **160**(3): 104–107.
- 14 Shinya K, Lanzetta M, Conolly WB. Risk and complications in endoscopic carpal tunnel release. *Journal of Hand Surgery—British Volume* 1995; **20**(2): 222–227.
- 15 Waegeneers S, Haentjens P, Wylock P. Operative treatment of carpal tunnel syndrome. *Acta Orthopaedica Belgica* 1993; **59**(4): 367–370.
- 16 Watson HK, Fong D. Dystrophy, recurrence, and salvage procedures in Dupuytren's contracture. *Hand Clinics* 1991; **7**(4): 745–755; discussion 57–58.
- 17 Prosser R, Conolly WB. Complications following surgical treatment for Dupuytren's contracture. *Journal of Hand Therapy* 1996; **9**(4): 344z–348.
- 18 Sampson SP, Wisch D, Badalamente MA. Complications of conservative and surgical treatment of de Quervain's disease and trigger fingers. *Hand Clinics* 1994; **10**(1): 73–82.
- 19 Mont MA, Antonaides S, Krackow KA, Hungerford DS. Total knee arthroplasty after failed high tibial osteotomy. A comparison with a matched group. *Clinical Orthopaedics and Related Research* 1994; **(299)**: 125–130.
- 20 Radin EL. Anterior tibial tubercle elevation in the young adult. *Orthopedic Clinics of North America* 1986; **17**(2): 297–302.
- 21 Katz MM, Hungerford DS, Krackow KA, Lennox DW. Reflex sympathetic dystrophy as a cause of poor results after total knee arthroplasty. *Journal of Arthroplasty* 1986; **1**(2): 117–124.
- 22 Myerson MS, McGarvey WC, Henderson MR, Hakim J. Morbidity after crush injuries to the foot. *Journal of Orthopaedic Trauma* 1994; **8**(4): 343–349.
- 23 Komenda GA, Myerson MS, Biddinger KR. Results of arthrodesis of the tarsometatarsal joints after traumatic injury. *Journal of Bone and Joint Surgery—American Volume* 1996; **78**(11): 1665–1676.
- 24 Isakov E, Susak Z, Korzets A. Reflex sympathetic dystrophy of the stump in below-knee amputees. *Clinical Journal of Pain* 1992; **8**(3): 270–275.
- 25 Mittal R, Khetarpal R, Malhotra R, Kumar R. The role of Tc-99m bone imaging in the management of pain after complicated total hip replacement. *Clinical Nuclear Medicine* 1997; **22**(9): 593–595.
- 26 Bach BR Jr, Wojtys EM, Lindenfeld TN. Reflex sympathetic dystrophy, patella infera contracture syndrome, and loss of motion following anterior cruciate ligament surgery. *Instructional Course Lectures* 1997; **46**: 251–260.
- 27 Anonymous. Complications of arthroscopy and arthroscopic surgery: results of national survey. *Committee on Complications of Arthroscopy Association of North America. Arthroscopy* 1985; **1**(4): 214–220.
- 28 Small NC. Complications in arthroscopic surgery performed by experienced arthroscopists. *Arthroscopy* 1988; **4**(3): 215–221.
- 29 Feder KS, Schonholtz GJ. Ankle arthroscopy: review and long-term results. *Foot and Ankle* 1992; **13**(7): 382–385.
- 30 Harrington P, Aiyaswami KV, Stephens MM. Diagnostic and therapeutic ankle arthroscopy. *Irish Journal of Medical Science* 1996; **165**(2): 121–124.
- 31 Petchkrua W, Weiss DJ, Patel RR. Reassessment of the incidence of complex regional pain syndrome type I following stroke. *Journal of Neurologic Rehabilitation* 2000; **14**(1): 59–63.
- 32 Geurts AC, Visschers BA, van Limbeek J, Ribbers GM. Systematic review of aetiology and treatment of post-stroke hand oedema and shoulder-hand syndrome. *Scandinavian Journal of Rehabilitation Medicine* 2000; **32**(1): 4–10.
- 33 Saddison DK, Vanek VW. Reflex sympathetic dystrophy after modified radical mastectomy: a case report (see comments). *Surgery* 1993; **114**(1): 116–120.
- 34 Poncelet C, Perdu M, Levy-Weil F, Philippe HJ, Nisand I. Reflex sympathetic dystrophy in pregnancy: nine cases and a review of the literature. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* 1999; **86**(1): 55–63.
- 35 Karras D, Karargiris G, Vassilopoulos D, Karatzetos C. Reflex sympathetic dystrophy syndrome and osteogenesis imperfecta. A report and review of the literature. *Journal of Rheumatology* 1993; **20**(1): 162–164.
- 36 Sherman OH, Fox JM, Snyder SJ, Del Pizzo W, Friedman MJ, Ferkel RD *et al.* Arthroscopy—'no-problem surgery'. An analysis of complications in two thousand six hundred and forty cases. *Journal of Bone and Joint Surgery—American Volume* 1986; **68**(2): 256–265.
- 37 Rauis AL. A series of 104 posttraumatic cases of reflex sympathetic dystrophy. *Acta Orthopaedica Belgica* 1999; **65**(1): 86–90.
- 38 Van Houdenhove B, Vervaeke G, Onghena P, Vasquez G, Vandeput C, Stans L *et al.* Psychometric characteristics of 66 patients with reflex sympathetic dystrophy. *European Journal of Pain* 1994; **15**(2–3): 50–58.
- 39 Van Houdenhove B. Neuro-algodystrophy: a psychiatrist's view. *Clinical Rheumatology* 1986; **5**(3): 399–406.
- 40 Geertzen JHB, De Bruijn-Kofman AT, De Bruijn HP, Van De Wiel HBM, Dijkstra PU. Stressful life events and psychological dysfunction in complex regional pain syndrome type I. *Clinical Journal of Pain* 1998; **14**(2): 143–147.

- 41 Geertzen JH, Dijkstra PU, Groothoff JW, ten Duis HJ, Eisma WH. Reflex sympathetic dystrophy of the upper extremity—a 5.5-year follow-up. Part II. Social life events, general health and changes in occupation. *Acta Orthopaedica Scandinavica Supplementum* 1998; **279**: 19–23.
- 42 Zucchini M, Alberti G, Moretti MP. Algodystrophy and related psychological features. *Functional Neurology* 1989; **4**(2): 153–156.
- 43 Feldman SI, Downey G, Schaffer-Neitz R. Pain, negative mood, and perceived support in chronic pain patients: a daily diary study of people with reflex sympathetic dystrophy syndrome. *Journal of Consulting & Clinical Psychology* 1999; **67**(5): 776–785.
- 44 Bruehl S, Carlson CR. Predisposing psychological factors in the development of reflex sympathetic dystrophy. A review of the empirical evidence. *Clinical Journal of Pain* 1992; **8**(4): 287–299.
- 45 Monti DA, Herring CL, Schwartzman RJ, Marchese M. Personality assessment of patients with complex regional pain syndrome type I. *Clinical Journal of Pain* 1998; **14**(4): 295–302.
- 46 Nelson DV, Novy DM. Psychological characteristics of reflex sympathetic dystrophy versus myofascial pain syndromes. *Regional Anesthesia* 1996; **21**(3): 202–208.
- 47 Van Houdenhove B, Vasquez G. Is there a relationship between reflex sympathetic dystrophy and helplessness? Case reports and a hypothesis. *General Hospital Psychiatry* 1993; **15**(5): 325–329.
- 48 Egle UT, Hoffmann SO. Psychosomatic connections in reflex sympathetic dystrophy or Sudeck's disease—survey of the literature and first clinical results. *Psychotherapie Psychosomatik Medizinische Psychologie* 1990; **40**(3–4): 123–135.
- 49 Birklein F, Riedl B, Sieweke N, Weber M, Neundorfer B. Neurological findings in complex regional pain syndrome—analysis of 145 cases. *Acta Neurologica Scandinavica* 2000; **101**(4): 262–269.
- 50 Verdugo RJ, Ochoa JL. Abnormal movements in complex regional pain syndrome: assessment of their nature. *Muscle and Nerve* 2000; **23**(2): 198–205.
- 51 Buijs EJ, Klijn FAM, Lindeman E, Van Wijck AJM. Reflex sympathetic dystrophy as a factitious disorder. *Nederlands Tijdschrift voor Geneeskunde* 2000; **144**(34): 1617–1620.
- 52 Neuschwander D, Drez D Jr, Heck S. Pain dysfunction syndrome of the knee. *Orthopedics* (Thorofare, NJ) 1996; **19**(1): 27–32.
- 53 Harden RN, Bruehl S, Galer BS, Saltz S, Bertram M, Backonja M *et al.* Complex regional pain syndrome: are the IASP diagnostic criteria valid and sufficiently comprehensive? *Pain* 1999; **83**(2): 211–219.
- 54 Cuartero-Plaza A, Martinez-Mirallas E, Benito-Ruiz P, Martinez-Pardo S, Sanz Marin MP. Abnormal bone scintigraphy and silent radiography in localized reflex sympathetic dystrophy syndrome. *European Journal of Nuclear Medicine* 1992; **19**(5): 330–333.
- 55 Todorovic-Tirnanic M, Obradovic V, Han R, Stankovic D, Sekulic D *et al.* Diagnostic approach to reflex sympathetic dystrophy after fracture: radiography or bone scintigraphy? *European Journal of Nuclear Medicine* 1995; **22**(10): 1187–1193.
- 56 Shahabpour M, Handelberg F, Opdecam M, Osteaux M, Spruyt D, Vaes P *et al.* Magnetic resonance imagine (MRI) of the ankle and hindfoot. *Acta Orthopaedica Belgica* 1992; **58**(suppl 1): 5–14.
- 57 Darbois H, Boyer B, Dubayle P, Lechevalier D, David H, Ait-Ameur A. MRI findings in reflex sympathetic dystrophy of the foot. *Journal de Radiologie* 1999; **80**(8): 849–854.
- 58 Graif M, Schweitzer ME, Marks B, Matteucci T, Mandel S. Synovial effusion in reflex sympathetic dystrophy: an additional sign for diagnosis and staging. *Skeletal Radiology* 1998; **27**(5): 262–265.
- 59 Gulevich SJ, Conwell TD, Lane J, Lockwood B, Schwettman RS, Rosenberg N *et al.* Stress infrared telethermography is useful in the diagnosis of complex regional pain syndrome, type I (formerly reflex sympathetic dystrophy). *Clinical Journal of Pain* 1997; **13**(1): 50–59.
- 60 Chapurlat RD, Duboeuf FP, Liens D, Meunier PJ. Dual energy X-ray absorptiometry in patients with lower limb reflex sympathetic dystrophy syndrome (see comments). *Journal of Rheumatology* 1996; **23**(9): 1557–1559.
- 61 Viel E, Ripart J, Pelissier JJ. Management of reflex sympathetic dystrophy. *Annales de Medecine Interne* 1999; **150**(3): 205–210.
- 62 Schuind F, Burny F. Can algodystrophy be prevented after hand surgery? *Hand Clinics* 1997; **13**(3): 455–476.
- 63 Gschwind C, Fricker R, Lacher G, Jung M. Does peri-operative guanethidine prevent reflex sympathetic dystrophy? *Journal of Hand Surgery—British Volume* 1995; **20**(6): 773–775.
- 64 Wang JK, Johnson KA, Ilstrup DM. Sympathetic blocks for reflex sympathetic dystrophy. *Pain* 1985; **23**(1): 13–17.
- 65 Severens JL, Oelermans HM, Weegels AJ *et al.* Cost-effectiveness analysis of adjuvant physical or occupational therapy for patients with reflex sympathetic dystrophy. *Archives of Physical Medicine and Rehabilitation* 1999; **80**(9): 1038–1043.
- 66 Vecht Ch J. Treatment of posttraumatic dystrophy. *Nederlands Tijdschrift voor Geneeskunde* 2000; **144**(34): 1620–1622.
- 67 Oerlemans HM, Oostendorp RAB, De Boo T, Van der Laan L, Severens JL, Goris RJA. Adjuvant physical therapy versus occupational therapy in patients with reflex sympathetic dystrophy/complex regional pain syndrome type I. *Archives of Physical Medicine and Rehabilitation* 2000; **81**(1): 49–56.
- 68 Zyluk A. The reasons for poor response to treatment of post-traumatic reflex sympathetic dystrophy. *Acta Orthopaedica Belgica* 1998; **64**(3): 309–313.
- 69 Kingery WS. A critical review of controlled clinical trials for peripheral neuropathic pain and complex regional pain syndromes (see comments). *Pain* 1997; **73**(2): 123–139.
- 70 Jadad AR, Carroll D, Glynn CJ, McQuay HJ. Intravenous regional sympathetic blockade for pain relief in reflex sympathetic dystrophy: a systematic review and a randomized, double-blind crossover study. *Journal of Pain and Symptom Management* 1995; **10**(1): 13–20.
- 71 Wechsler RJ, Frank ED, Halpern EH, Nazarian LN, Jalali S, Ratner ER. Percutaneous lumbar sympathetic plexus catheter placement for short- and long-term pain relief: CT technique and results. *Journal of Computer Assisted Tomography* 1998; **22**(4): 518–523.
- 72 Katz MM, Hungerford DS. Reflex sympathetic dystrophy affecting the knee. *Journal of Bone and Joint Surgery—British Volume* 1987; **69**(5): 797–803.
- 73 Urschel HC Jr. Dorsal sympathectomy and management of thoracic outlet syndrome with VATS. *Annals of Thoracic Surgery* 1993; **56**(3): 717–720.
- 74 Lacroix H, Vander Velpen G, Penninckx F, Nevelsteen A, Suy R. Technique and early results of videoscopic lumbar sympathectomy. *Acta Chirurgica Belgica* 1996; **96**(1): 11–14.
- 75 Rocco AG. Radiofrequency lumbar sympathectomy. The evolution of a technique for managing sympathetically maintained pain. *Regional Anesthesia* 1995; **20**(1): 3–12.
- 76 Cortet B, Flipo RM, Coquerelle P, Duquesnoy B, Delcambre B. Treatment of severe, recalcitrant reflex sympathetic dystrophy: assessment of efficacy and safety of the second generation bisphosphonate pamidronate. *Clinical Rheumatology* 1997; **16**(1): 51–56.
- 77 Maresca V. Human calcitonin in the management of osteoporosis: a multicentre study. *Journal of International Medical Research* 1985; **13**(6): 311–316.
- 78 Becker WJ, Ablett DP, Harris CJ, Dold ON. Long term treatment of intractable reflex sympathetic dystrophy with intrathecal morphine (see comments). *Canadian Journal of Neurological Sciences* 1995; **22**(2): 153–159.

- 79 Tountas AA, Noguchi A. Treatment of posttraumatic reflex sympathetic dystrophy syndrome (RSDS) with intravenous blocks of a mixture of corticosteroid and lidocaine: a retrospective review of 17 consecutive cases. *Journal of Orthopaedic Trauma* 1991; **5**(4): 412–419.
- 80 Grundberg AB. Reflex sympathetic dystrophy: treatment with long-acting intramuscular corticosteroids. *Journal of Hand Surgery American* 1996; **21**(4): 667–670.
- 81 Kumar K, Nath RK, Toth C. Spinal cord stimulation is effective in the management of reflex sympathetic dystrophy. *Neurosurgery* 1997; **40**(3): 503–508; discussion 8–9.
- 82 Kemler MA, Barendse GAM, Van Kleef M, De Vet HCW, Rijks CPM, Furnee CA *et al.* Spinal cord stimulation in patients with chronic reflex sympathetic dystrophy. *New England Journal of Medicine* 2000; **343**(9): 618–624.
- 83 Seale KS. Reflex sympathetic dystrophy of the lower extremity. *Clinical Orthopaedics & Related Research* 1989; **243**: 80–85.
- 84 Ochoa JL, Verdugo RJ. Reflex sympathetic dystrophy. A common clinical avenue for somatoform expression. *Neurologic Clinics* 1995; **13**(2): 351–363.
- 85 Cooney WP. Somatic versus sympathetic mediated chronic limb pain. Experience and treatment options. *Hand Clinics* 1997; **13**(3): 355–361.
- 86 Harden RN, Cole PA. New developments in rehabilitation of neuropathic pain syndromes. *Neurologic Clinics* 1998; **16**(4): 937–950.
- 87 Stanton-Hicks M, Baron R, Boas R, Gordh T, Harden N, Hendler N *et al.* Complex regional pain syndromes: guidelines for therapy (see comments). *Clinical Journal of Pain* 1998; **14**(2): 155–166.
- 88 Harden RN. A clinical approach to complex regional pain syndrome. *Clinical Journal of Pain* 2000; **16**(2 suppl): S26–S32.
- 89 Muller LP, Muller LA, Happ J, Kerschbaumer F. Frozen shoulder: a sympathetic dystrophy? *Archives of Orthopaedic and Trauma Surgery* 2000; **120**(1–2): 84–87.
- 90 Julian TM, Backonja MM. Vulvar vestibulis syndrome: a form of reflex sympathetic dystrophy? *Journal of Lower Genital Tract Disease* 2000; **4**(1): 26–29.
- 91 Reading AE. A critical analysis of psychological factors in the management and treatment of chronic pelvic pain. *International Journal of Psychiatry in Medicine* 1982; **12**(2): 129–139.
- 92 Robson S, Henshaw R. Intractable pelvic pain following Filshie clip application. *Australian and New Zealand Journal of Obstetrics and Gynaecology* 1997; **37**(2): 242–243.
- 93 Gross RJ, Doerr H, Caldirola D, Guzinski GM, Ripley HS. Borderline syndrome and incest in chronic pelvic pain patients. *International Journal of Psychiatry in Medicine* 1980; **10**(1): 79–96.
- 94 Choa RG, Swami KS. Testicular denervation: a new surgical procedure for intractable testicular pain. *British Journal of Urology* 1992; **70**: 417–419.
- 95 Cadeddu JA, Bishoff JT, Chan DY, Moore RG, Kavoussi LR, Jarrett TW. Laparoscopic testicular denervation for chronic orchalgia. *The Journal of Urology* 1999; **162**: 733–736.